Elephantopus scaber Linn.

Family : Asteraceae

Parts Used: Root, Leaf, Flower

Vernacular Names

English : Prickly leaved elephant's foot

Malayalam : Anachuvadi
Hindi : Gobhi
Sanskrit : Hastipadi
Bengali : Gajilata
Gujarathi : Bhopatahari
Kannada : Nayee nalige
Tamil : Anashovadi



Distribution and Habitat: Found throughout India, especially in dry localities.

Botany: A rigid herb, 45 cm tall, usually perennial with short root stock; stem dichotomously branched

- *Leaves:* Obovate- oblong, mostly radical forming a spreading rosette on the ground, hairy on both surfaces, 12.5- 20 cm long and 3.8-5.7 cm broad
- Flowers: Purple in heads, heads numerous, closely packed
- Fruits: Achene, pappus of 4-6 bristles dilated at the base.

Properties: Astringent, febrifuge

Chemical constituents: Epifriedeliniol, lupeol, stigmasterol and a mixture of triacontan-1-ol and dotriaconta-1-ol. Sequiterpene dilactone. Alcoholic extract contains antibiotic elephentopin

Uses: It is used in dysuria, intermittent fevers, bronchitis, hemorrhoids and hepatopathy, ophthalmopathy, swellings, cough and skin diseases





