

Tabernaemontana dichotoma L.

Synonyms : *Tabernaemontana alternifolia* L.

Family : Apocynaceae

Parts Used : Root , Flower, Latex

Vernacular Names

English : *Tabernaemontana*

Malayalam : Koonampala



Botany: A small tree; branches very stout, woody, cylindric; buds resinous.

• **Leaves:** 10-20 X 2.5-5.0 cm, pale yellow brown when dry, midrib very stout, nerves 16-22 pairs; petiole 0.6-1.8 cm, base semiamplexicaul.

• **Flowers** : Peduncle 5.0-12.5 cm, bracts nil. Calyx hemispheric, very thick, lobes semi circular. Flowers scarcely fragrant. Corolla white, tube 1.8-2.5 cm, yellow, cylindric, swollen at the very base, limb white, 5-7.5 cm diameter, lobes broadly oblong, falcate, obtuse. Anthers half way down the tube. Follicles usually solitary, 10-12.5 X 2.5-5.0 cm, recurved, oblong, narrowed at both ends, dorsally concave, ventrally 2-ribbed, orange yellow.

• **Seeds** : 1.8 cm long, buried in scarlet pulp.

Properties: Anti-inflammatory, opthalmic, antidiarrhoeal, vermicide

Chemical constituents: tabernamine, 16,22-dihydro-16-hydroxyapparicine, dichomine, vallesamine, voacamine

Uses: Root juice for eye infections. crushed bark and leaves used as antiseptic on wounds and for treating ulcers and fistula.

Propagation: Cuttings

